

Biogeochemistry with many tracers at high resolution




Positive Definite Redi

- Biogeochem may require positive definite advection/diffusion (nonlinear equations sensitive to over/undershoots)
- Interest in marginal species
- Steep gradients:



Positive Definite Redi

- Biogeochem may require positive definite advection/diffusion (nonlinear equations sensitive to over/undershoots)
- Interest in marginal species
- Steep gradients:



Positive Definite Redi


- Biogeochem may require positive definite advection/diffusion (nonlinear equations sensitive to over/undershoots)
- Interest in marginal species

Steep gradients:
 Scale outgoing flux to less than tracer content of cell [Bott/Smolarkiewicz]
 Used for aquaplanet and phytoplankton mutations

Many passive tracers

Added to MITgcm kernel:

- Full support for ~3000 tracers (incl. diagnostics)
 [Chris Hill]
- Long tracer time step (pkg/longstep)



Makes possible:


- 78 species (99 ptracers)
- cs510 grid (1/6° resolution)

at 3-4 months/day on 360 cores.

Many passive tracers

Added to MITgcm kernel:

- Full support for ~3000 tracers (incl. diagnostics)
 [Chris Hill]
- Long tracer time step (pkg/longstep)



Makes possible:

- 78 species (99 ptracers)
- cs510 grid (1/6° resolution)

at 3-4 months/day on 360 cores.

DO_STATEVARS_DIAGS 16.85s

DO_OCEANIC_PHYS

DYNAMICS 33.64s

SOLVE FOR PRESSURE 13.27s

BLOCKING_EXCHANGES 51.54s

THERMODYNAMICS
33.8s

ptracer adv/diff

LONGSTEP_THERMODYNAMICS

bio eq.

GCHEM_FORCING_SEP 100.7s